GAT Grupo Português de Activistas sobre Tratamentos de VIH/SIDA Pedro Santos



Pilot HCV rapid testing project among MSM Checklist project, Lisbon, Portugal

<u>M. Rocha¹</u>, M.J. Campos¹, R. Fuertes¹, J. Cornejo¹, J. Brito¹, J. Esteves¹, F. Ferreira¹, H. Machado¹, N. Pinto¹, L. Mendão¹


¹GAT Portugal | CheckpointLX

HCV prevalence Europe

- 7,9/100 000 persons
- 25 34 years (28,2% cases)
- Male-to-female ratio of 1.9:1
- Acute HCV cases:
 - 33,3 % PWID
 - 24,4% MSM











HCV incidence and prevalence HepH Portugal Insufficient data for prevalence estimate

 Incidence of 1/100 000 persons per year and prevalence of 1-1,5% estimates (Anjo *et al*, 2014)


(ECDCb, 2010; ECDCc,





2010);









Targeted HCV screening Specific criteria



- Sharing:
 - drug preparation or injecting equipment;
 - drug-snorting paraphernalia;
- Unprotected anal group sex;
- Group sex without using new condoms between partners;
- Insertive or receptive fisting;
- Having had sex with presence of blood.





CheckpointLX

Peer-led HIV community based anonymous counseling testing and linkage center for MSM

- Targeted HCV screening strategy
 - Anonymity
 - Observance of at least one of the screening criteria
 - Report without disclosure
 - Both HIV negative or positive were eligible
- Referral to Checklist project (*express link*)
 - MSM medical team granting STI screenings
 - Link to infectious disease clinic to confirmatory the results








HCV targeted screening Rapid test results









HCV targeted screening Pilot study findings



- Peer-led HIV CBACTL center are na opportunistic setting for targeted HCV screenings
 - HCV targeted screening are effective here
- Enabling strategies for MSM adherence to HCV screenings
 - Anonymity
 - Adding specific high-risk sexual practices of MSM
 - Choice of criteria report without disclosure/ specifying the behavior
- HCV targeted screening to MSM should a standard practice at HIV CBACTL centers






HCV targeted screening 2014 new criteria and updates

- Poppers bottle included in the drug-snorting paraphernalia
- Shared tattoo or piercing puncture paraphernalia
- Person living with HIV that had any of the criteria mentioned since last follow-up/ screening at infectious desease clinic










HCV targeted screening Aug 2014 to Sep 2014









GAT Grupo Português de Activistas sobre Tratamentos de VIH/SIDA Pedro Santos



Pilot HCV rapid testing project among MSM Checklist project, Lisbon, Portugal

<u>M. Rocha¹</u>, M.J. Campos¹, R. Fuertes¹, J. Cornejo¹, J. Brito¹, J. Esteves¹, F. Ferreira¹, H. Machado¹, N. Pinto¹, L. Mendão¹

¹GAT Portugal | CheckpointLX

