# HIV and viral hepatitis response under UHC approach – WHO guidance on testing strategies and linkage to care

#### Elena Vovc, MD, MPH

Joint TB, HIV and viral hepatitis programme
Division of Health Emergencies & Communicable Diseases
WHO Regional Office for Europe, Copenhagen
vovce@who.int













#### Late diagnosis – key challenge across the Region

% of people testing positive for HIV who are diagnosed late (CD4<350) by age range













#### Newer strategies: reaching the unreached

- Reduce time of diagnosis confirmation:
  - Simplified testing strategies
  - Rapid diagnostic technologies
- Tailor testing strategies to the context of the epidemic
- Trained lay providers can safely and effectively perform HIV testing services
- Since 2016, WHO recommends self-testing as an additional option to encourage HIV diagnosis
- Voluntary assisted partner notification as part of a comprehensive package of testing and care for PLHIV













# Evolution of HTC Approaches

1985
1990 VCT sites, e.g. AIC Uganda

1985
Clinical diagnostic testing;
Blood donors
1999 Social marketing, e.g. New Start

2000 Campaigns
2007 PITC (WHO)





2013 Community-based HTC (WHO)



2015






- Confirm HIV test results in 14 days
- Put on ART in **7 days**

#### **Testing strategies**

#### High prevalence (>5%)












#### Test for triage

- Most useful for
  - HIV self-testing
  - Community-based testing
  - Settings where all three assays can not be conducted reliably




#### Not useful for

Same-day diagnosis









#### WHO HTS Info: new app







http://www.who.int/hiv/mediacentre/news/hts-info-app/en/











## HIV testing approaches in the WHO European Region **2016**



#### HIV testing approches in central and eastern Europe, 2017













# Countries change testing policies for better impact

- RDT use, decentralized services - reduce time of test confirmation results
- Community based testing great impact at country level
- Integrated testing for hepatitis and expanding testing sites brings good national level results

Berlin, 22-23 January 2019 WHO Regional workshop on HIV and hepatitis testing and PrEP






ная организация охранения

кое региональное бюро

#### 2018 HIVST policy and implementation

















#### Regional essential targets by 2020 towards elimination of hepatitis



# GUIDELINES ON HEPATITIS B AND C TESTING FEBRUARY 2017

#### ■ 2015 BASELINE



<sup>\*</sup>Measuring the progress on vertical transmission prevention is limited by data on pregnant women screening coverage

PMTCT – prevention of mother-to-child transmission









<sup>\*\*</sup> Measuring the progress on HBV treatment is now limited by the absence of data on the proportion of persons eligible

Diagnosis of chronic HCV infection



Diagnosis of chronic HBV infection




#### Linkage to care

- Linkage a process of actions and activities that supports people testing for HIV and people diagnosed with HIV in engaging with prevention, treatment and care services as appropriate for their HIV status.
- For people with HIV, it refers to the period beginning with HIV diagnosis and ending with enrolment in care or treatment













## Global indicators for the monitoring and evaluation of the health sector response to HIV



#### Achieving 90-90-90 and SDG 3.3

Universal Health Coverage: is it possible in the context of HIV epidemic in the WHO European Region?



universal health coverage, the continuum of services, a public health approach













### HIV resource availability 2006-2016 and projected resource needs by 2020, Eastern Europe and Central Asia (EECA)



#### UHC – a people-centered approach











# Making UHC work for the HIV Epidemic in our Region

- **Effective interventions to be brought to scale** and included in UHC packages (simplify strategies)
- Innovations are needed to increase efficiencies and sustainability
- People-centered services and differentiated service delivery (decentralize HTS)
- Services for key populations ending AIDS not possible without addressing the epidemics within key populations and their partners
- Advocate for increased health budgets and strategic funding allocations for comprehensive HIV services as part of UHC package
- Invest in community services to reach out to those in need and address HIV-related stigma
- A shared challenge power through partnership










#### Acknowledgments:

Masoud Dara
Antons Mozalevskis
Anita Sands
Cheryl Johnson

#### WHO key resources



- http://www.who.int/hiv/pub/guidelines/hiv-testing-services/en/
- http://www.who.int/hiv/pub/self-testing/hiv-self-testing-guidelines/en/
- https://www.who.int/hepatitis/publications/guidelines-hepatitis-c-b-testing/en/







